

A LOWER DEVONIAN (EMSIAN) SPECIES OF THE GENUS *Bactrocrinites* (CRINOIDEA): *Bactrocrinites robustus* n. sp. (CANTABRIAN MOUNTAINS, NW SPAIN)

Raimundo PIDAL

Departamento de Geología, Universidad de Oviedo; Jesús Arias de Velasco, s/n. 33005 Oviedo. e-mail: pidal@ozu.es

Pidal, R. 2008. A Lower Devonian (Emsian) species of the genus *Bactrocrinites* (Crinoidea): *Bactrocrinites robustus* n. sp. (Cantabrian Mountains, NW Spain). [Una especie del Devónico Inferior (Emsiense) del género *Bactrocrinites* (Crinoidea): *Bactrocrinites robustus* n. sp. (Cordillera Cantábrica, NO de España).] *Revista Española de Paleontología*, **23** (2), 267-271. ISSN 0213-6937.

ABSTRACT

The only previous reference to Spanish bactrocrinitids (Crinoids) is found in Breimer (1962), who reports a specimen catalogued as TB73 on show at the museum of the Instituto Geológico y Minero de España (IGME) in Madrid. This specimen was found in the locality of Colle, León (Cantabrian Mountains) in rocks of the La Vid Group (Emsian). It was classified as *Bactrocrinites* sp., and was considered to be related to *B. fusiformis* (Roemer, 1844) and to *B. muelleri* (Jaekel, 1895). New and exhaustive search in the locality of Colle have led to the discovery of two new specimens of the genus whose characteristics allow a new species to be defined: *B. robustus* n. sp., with a much more robust calyx than the other known species of the genus.

Keywords: Crinoidea, Inadunata, *Bactrocrinites*, Emsian, La Vid Group, Cantabrian Mountains, Spain.

RESUMEN

La única referencia previa a los bactrocrinos (Crinoideos) españoles se debe a Breimer (1962) y se refiere a un espécimen expuesto en el museo del Instituto Geológico y Minero de España (IGME) catalogado como TB73. Dicho espécimen fue recogido en la localidad leonesa de Colle (Cordillera Cantábrica) en rocas del Grupo La Vid (Emsiense). El ejemplar fue clasificado como *Bactrocrinites* sp., y considerado afín a *B. fusiformis* (Roemer, 1844) y a *B. muelleri* (Jaekel, 1895). Nuevas campañas de búsqueda exhaustiva en la localidad de Colle culminaron con el hallazgo de dos nuevos ejemplares del género cuyas características permiten definir una nueva especie: *B. robustus* n. sp., con cáliz mucho más masivo que el de las demás especies conocidas del género.

Palabras clave: Crinoideos, Inadunata, *Bactrocrinites*, Emsiense, Grupo La Vid, Cordillera Cantábrica, España.

INTRODUCTION

To date, the first and only reference of bactrocrinitids in Spain is to be found in the famous monograph on Spanish crinoids by A. Breimer (1962). This study reports the existence of a specimen catalogued as TB73, from the Emsian period of the La Vid Formation in the locality of Colle, León (Fig. 1), Level Crin 1 (probably equal to Interval 27 of García-Alcalde, 1996; Fig. 2), stored in the collection of the Instituto Geológico y Minero de España (IGME), in Madrid. Breimer described and classified this example as *Bactrocrinites* sp., pointing to the possibility of its being a

new species, related to *B. fusiformis* (Roemer, 1844) or to *B. muelleri* (Jaekel, 1895). The discovery of additional examples of the genus at the same locality in León, possibly from the same level as Breimer's specimen, enable us to both broaden and refine our knowledge of Spanish bactrocrinitids via the description of a new species, *Bactrocrinites robustus* n. sp. From a palaeoecological viewpoint, the individuals of the species, which are completely benthonic, must have lived in a shallow, neritic environment, in keeping with the characteristics of the rocks in which they are to be found (Álvarez, 1990; Álvarez & Brime, 1982; García-Alcalde, 1996, 1998; García-Alcalde *et al.*, 2002).



Figure 1. Geographical and Geological situation of Collé (from García-Alcalde, 1998).

The material of the new species is stored at the Museum of the Department of Geology (Section Paleontology) (DPO) of the Oviedo University (Asturias, Spain).

SYSTEMATIC PALAEOONTOLOGY

Class CRINOIDEA Miller, 1821

Subclass INADUNATA Moore & Laudon, 1943

Suborder CYATHOCRININA Bather, 1899

Family **Bactrocrinitidae** Jaekel, 1918

Genus *Bactrocrinites* Schnur, 1849

Type species: *Bactrocrinites fusiformis* Roemer, 1844.

Bactrocrinites robustus n. sp.

Fig. 3. Table 1

Derivatio nominis: Indicates the robust, thick calyx of the type material, an unusual peculiarity in *Bactrocrinites*.

Material: The Holotype, DPO 113601, and another fragmentary example, Paratype DPO 113602, from the locus and stratum typicum. Both with evident signs of dissolution.

Locus and stratum typicum: Both the holotype as well as the paratype come from argillaceous limestones from Level 27 (Fig. 2), possibly coincident with Breimer’s CRIN 1 (1962), cropping out in the locality of Collé, in León (Fig. 1) in the La Vid Group (Emsian).

Diagnosis: Dicyclic calyx, with a conical-globular longitudinal profile, somewhat flattened at the base; robust, highly convex plates; appreciably reduced radial with respect to the anal X; very strong arms and a relatively thin stem.




Figure 2. Stratigraphic column of the Lower Devonian at Collé, indicating Level 27, characterized by abundant crinoid and blastoid fauna, the locus and stratum typicum of *Bactrocrinites robustus* n. sp. (modified from García-Alcalde, 1998)

Description based mainly on the holotype: Dicyclic calyx, with a conical-globular longitudinal profile and approximate dimensions: height, 23 mm and maximum width, 23 mm, made up of large, robust, convex plates, including the radianals. The (incomplete) tegmen appears to possess a reduced number of plates. No ornamentation is observed (Fig. 3 a, b, c, d, e and Table 1).

Infrabasal circle is made up of five equally sized pentagonal plates, forming a small concavity at the base, just in the area of the connection with the stem. These plates respectively measure 7 and 6.5 mm along the mean vertical axis (MVA) and mean transversal axis (MTA). Aborally, the infrabasal set presents a weakly five-lobed contour. The anteroposterior axis (APA) measures 8 mm. The set limits distally with the basal plates. Circular stem scar with a diameter of 3.5 mm. Axial canal is not conserved.

Five basal plates: two heptagonal, the posterior and right posterior (the seventh side being developed to better accommodate the anal area) and three hexagonal. The plates measure 10 and 8.5 mm, respectively, along the usual axes. The right posterior basal plate bears the radial, and the posterior the radial and the anal X.




Figure 3. Organization of the calyx of *Bactrocrinites robustus* n. sp. a) holotype DPO 113601, posterior view, radial in black, and b) anterior view (camera lucida drawings); c) holotype DPO 113601, anterior view; d) posterior view; e) aboral view; f) paratype DPO 113602, showing one radial and seven contiguous brachial plates.

The radial cycle is composed of five plates: four pentagonal, and the somewhat smaller right posterior, which is hexagonal. The upper side of all the plates is markedly concave. The pentagonal plates measure 7 and 10 mm, respectively, along the MVA and MTA, and the hexagonal, 5 and 10 mm. The radial cycle limits proximally with the basal, but the right posterior radial also does so with the radianal and with the anal X, and the left posterior with the anal X. They are distally arranged in net contact with the brachial series.

The beginning of an anal area is observed in the calyx, made up of two plates of different sizes: the radianal and the anal X. The former, a typical quadrangle, with sides measuring 3 mm, is obliquely arranged with respect to the right posterior radial and limits proximally with two basal plates, the posterior and right posterior, and distally with the aforementioned radial and with the anal X. The anal X is an irregular pentagon and is considerably larger than the radianal, measuring 7 mm along the mean vertical axis; it is laterally in contact with the right posterior and left posterior radials, and rests proximally on the posterior basal and the radianal. The anal series appears to continue in the tegmen, where an additional plate can be observed that is longer than it is wide, 4 and 2 mm, respectively.

The tegmen is poorly preserved, not allowing a reliable description.

Calyx with five arms, of which only the first plates are conserved. These are equidimensional and strong, and are wider than high, 7 and 3 mm, respectively; its transversal section is roughly horseshoe in shape. The food groove is narrow and extends until almost the centre of the plates. Crenulation is not observed on the articular surface. The thick arms occupy practically all the oral area, and so it appears unlikely that this was made up of a large number of plates.

A maximum of only four primibrachials are conserved, in some series, in the holotype. Thus the total length of the arms is unknown, as is the answer to the question as to whether they branched or not. In the paratype, DPO 113602 (Fig. 3 f and Table 1), only two pentagonal radials are observed, measuring 8 and 11 mm, respectively (Table I), along the usual axes, and a series of seven primibrachials continuing on from one of the radials, which are wider than high, that respectively measure 7 and 3 mm.

Among the material adhering to the holotype, two kind of pieces have been identified: brachial and columnar, both in a poor state of conservation, resulting in their description being tentative. The brachial pieces appear to be characteristic of the ends of the arms, in this case, *B. robustus* n. sp. would possess considerably long brachial appendices. The pieces identified as columnar are thin. Their diameter varies between 3 and 3.5 mm, or even less. These dimensions coincide with those of the basal insertion. The columnar pieces have a circular to slightly five-lobed contour. The crenularium, which is petaloid in appearance, is perpendicular to the external edge of the columnar and occupies a major part of the articular facet; the areola and axial canal are star-shaped, whereas the articulation appears to be of the "simplexy" type.

To indicate that, though the above mentioned material seems to belong to the holotype, we cannot assure it for the present time. It will be necessary to hope that future finds confirm it.

No ornamentation is observed.

Discussion: Provisionally, the classificatory scheme adopted is that of Moore and Teichert in the "Treatise on Inver-

Specimen	Total height	Maximum width	Index Th/Mw	IBB MVA MTA	BB MVA MTA	RR MVA MTA	Brr Width and Height	Radial	Anal X
<i>Bactrocrinites</i> sp. TB73 of Breimer	22	15	1.47	9 5	8.5 6	4.5 6.5		4	4
<i>Bactrocrinites robustus</i> n. sp. 113601 Holotype	23	23	1	7 6.5	10 8.5	7 10	7 3	3	7
113602 Paratype						8 11	7 3		

Table 1. Measurements (in mm) of the *Bactrocrinites* specimens found in Colle (León), La Vid Group, Emsian. Th/Mw, total height/maximum width and MVA and MTA, mean vertical axis and mean transversal axis, respectively.

tebrate Paleontology” (1978). In fact, although Simms & Sevastopulo (1993) and Ausich (1998) proposed substantial changes in such scheme, many doubts continue to exist with respect to bactrocrinitids (see, for example, McIntosh, 1979, and McIntosh & Brett, 1988). We have therefore chosen to maintain the old scheme of the “Treatise”.

The attribution of a new species to the genus *Bactrocrinites* is justified by the existence of morphological affinities with the type species *B. fusiformis*: a tall conical dorsal cup, two anal plates in the cup: a quadrangular radial and a pentagonal anal X, the latter in line with the radials; radials that are significantly wider than high; basals that are higher than wide; infrabasals that are always laterally visible, generally higher than wide; etc. However, *B. robustus* is differentiated from *B. fusiformis* by the presence of a globular calyx, as high as wide, with prominent, markedly convex plates that confer it a characteristically robust appearance, in clear contrast to that of *B. fusiformis*, much more graceful and slim, made up of smooth plates that are barely convex.

Another important distinctive characteristic of *B. robustus* n. sp. is the presence in the calyx of a radial that is unusually smaller than the anal X. This aspect justifies the separation of the new species not only from the type species of the genus, but also from all the known forms of *Bactrocrinites*.

In particular, the new species is separated from the form classified by Breimer (1962) as *Bactrocrinites* sp. in the possession of a globular calyx, as tall as wide, made up of markedly convex plates. This aspect contrasts especially with that of Breimer’s very slim specimen, longer than wide and with flattened plates.

As regards the radial plates, those of *B. robustus* expand noticeably, while those of *Bactrocrinites* sp. incline inward. Likewise, those of *B. robustus* show the equal values as the axis MTA, both at the top level as in de lower, while in Breimer’s specimen the maximum width is at the lower level. As to the radial and anal X, *B. robustus* presents a much smaller radial than anal X. This contrasts not only with what occurs in *Bactrocrinites* sp., where the measurements of both plates are similar, but also with the rest of the known species of *Bactrocrinites*.

In keeping with its calyx, the arms of *B. robustus* n. sp. are powerful, whereas in Breimer’s specimen, according to the conserved primibrachial, they must have been more delicate.

The general characteristics of Breimer’s specimen situate it closer above all to *Bactrocrinites muelleri*.

ACKNOWLEDGEMENTS

I wish to express my gratitude to Jenaro García-Alcalde, Professor of Paleontology at the Geology Department of the University of Oviedo, as well as to the members and collaborators of this department (as is the case of the photographer, Joaquín Vázquez) for their help and advice given during the development of this work. A special mention to Professor Jean Le Menn, of the Paleontology Department of the University of Brest, who not only carried out the final review of this manuscript, but also contributed ideas to it, at the same time as well as facilitating oral information and abundant bibliographical material.

This work is a contribution to the University of Oviedo project: “Origin of Asturian Paleontology Fauna in the area of Ferroñes” and it has been financed by de project MEC- 05 - CGL 2005 - 03715 “Fases de desarrollo de asociaciones de corales y estromatoporoides en el Devónico Inferior de la Cordillera Cantábrica (NO de España).

REFERENCES

- Álvarez, F. 1990. Devonian Athyrid brachiopods from the Cantabrian Zone (NW Spain). *Biostratigraphie du Paléozoïque*, **11**, 1-311.
- Álvarez, F. & Brime, C. 1982. Aportaciones al conocimiento de las condiciones de formación de algunos depósitos fosilíferos del Devónico Cantábrico. *Trabajos de Geología, Universidad de Oviedo*, **12**, 153-157.
- Ausich, W. I. 1998. Early Phylogeny and Subclass Division of the Crinoidea (Phylum Echinodermata). *Journal of Paleontology*, **72**, 499-510.
- Ausich, W. I. & Kammer, T. W. 2001. The Study of Crinoids during the 20th Century and the Challenges of the 21st Century. *Journal of Paleontology*, **75**, 1161-1173.

- Breimer, A. 1962. A monograph on Spanish Palaeozoic Crinoidea. *Leidse Geologische Mededelingen*, **27**, 1-190.
- García-Alcalde, J. L. 1996. El Devónico del dominio Astur-Leonés en la Zona Cantábrica (N de España). *Revista Española de Paleontología*, n° **extraordinario**, 78-51.
- García-Alcalde, J. L. 1998. Terebratúlidos (Braquiópodos) del Emsiense superior de Colle (Sabero, León, N. de España). *Trabajos de Geología, Universidad de Oviedo*, **21**, 159-176.
- García-Alcalde, J. L., Carls, P., Pardo Alonso, M. V., Sanz López, J., Soto, F., Truyols-Massoni, M. & Valenzuela-Ríos, J. I. 2002. Devonian. In: *The Geology of Spain* (Eds. W. Gibbons & T. Moreno). The Geological Society of London, 67-91.
- Jaekel, O. 1895. Beiträge zur Kenntniss der palaeozoischen Crinoiden Deutschlands. *Pälaeontologischen Abhandlungen*, N. F., **3**, 1-116.
- Le Menn, J. 1985. Les Crinoïds du Dévonien Inferieur et Moyen du Massif Armoricaire. *Mémoires de la Société Géologique et Minéralogique de Bretagne*, **30**, 1-268.
- Le Menn, J. 1987. Growth Patterns and Evolutionary Trends of Devonian Crinoid Columns. *Geobios*, **20**, 811-829.
- McIntosh, G. C. 1979. Abnormal specimens of the Middle Devonian Crinoid *Bactrocrinites* and their effect on the taxonomy of the genus. *Journal of Paleontology*, **53**, 18-28.
- McIntosh, G. C. & Brett, C. E. 1988. Occurrence of the Cladid Inadunate Crinoid *Thalamocrinus* in the Silurian (Wenlockian of New York and Ontario). *Life Sciences Contributions*, **149**, 1-17.
- Méndez-Bedia, I. & Soto, F. 1984. Paleocological succession in a Devonian organic buildup (Moniello Fm., Cantabrian Mountains, NW Spain). *Geobios*, **8**, 151-157.
- Miller, S. A. & Gurley, W. F. E. 1895. New and interesting species of Paleozoic fossils. *Bulletin of the Illinois State Museum of Natural History*, **7**, 1-89.
- Moore, R. C., Jeffords, R. M. & Miller, T. H. 1968. Morphological Features of Crinoid Columns. *Echinodermata*, **8** (45), 1-30.
- Ubaghs, G. 1978. Classification of the Echinoderms. In: *Treatise on Invertebrate Palaeontology. Pt. T. Echinodermata 2.* (Eds. R.C. Moore & C. Teichert). The Geological Society of America and The University of Kansas, 359-401.
- Reed, F. R. C. 1908. Devonian faunas of the Northern Shan states. *Indian Geological Survey, Memoirs*, **2** (5), 1-157.
- Roemer, C. F. 1844. *Das Rheinische Uebergangsgebirge. Eine palaeontologisch-geognostische Darstellung.* Hanovre (Hahn), 96 pp.
- Schmidt, W. E. 1934. Die Crinoideen des Rheinischen Devons. Teil I: Die Crinoideen des Hunsrückschiefer. *Abhandlungen Preussischen Geologisches Landesanstalt*, N. F., **163**, 1-149.
- Schmidt, W. E. 1941. Die Crinoideen des Rheinischen Devons. Teil II, A. Nachtrag zu: Die Crinoideen des Hunsrückschiefers bis zur *Cultrijugatus*-Zone (mit Ausschluss des Hunsrückschiefers). *Abhandlungen Reichsstelle Bodenforschung*, N. F., **182**, 1-253.
- Steininger, J. 1849. Die Versteinerungen Des Uebergangsgebirges der Eifel. *Jahresbeiricht über den Schul-Cursus 1848-49 an dem Gymnasium zu Trier*, 1-50.
- Sieverts-Doreck, H. 1950. Über *Hexacrinus* und *Bactrocrinus*. *Neues Jahrbuch Geologisches Paläontologisches Monatsheften*, **3**, 80-87.
- Simms, M. J. & Sevastopulo, G. D. 1993. The origin of articulate crinoids. *Paleontology*, **36**, 91-109.

Manuscrito recibido: 18 de Febrero, 2008

Manuscrito aceptado: 20 de Septiembre, 2008

